

Physiological carbon dioxide, bicarbonate, and pH sensing

Martin Tresguerres · Jochen Buck · Lonny R. Levin

Received: 2 July 2010 / Revised: 16 July 2010 / Accepted: 18 July 2010 / Published online: 4 August 2010
© Springer-Verlag 2010

Abstract In biological systems, carbon dioxide exists in equilibrium with bicarbonate and protons. The individual components of this equilibrium (i.e., CO_2 , HCO_3^- , and H^+), which must be sensed to be able to maintain cellular and organismal pH, also function as signals to modulate multiple physiological functions. Yet, the molecular sensors for $\text{CO}_2/\text{HCO}_3^-/\text{pH}$ remained unknown until recently. Here, we review recent progress in delineating molecular and cellular mechanisms for sensing CO_2 , HCO_3^- , and pH.

Keywords Adenylyl cyclase · Bicarbonate · Carbon dioxide · Carbonic anhydrase · Channels · pH · Protons · Sensory transduction

Carbon dioxide (CO_2), bicarbonate ions (HCO_3^-), and pH/protons (H^+) are inextricably linked in biological systems. Due to ubiquitous carbonic anhydrases, CO_2 is in nearly instantaneous equilibrium with its hydrated form H_2CO_3 , which in turn rapidly dissociates into H^+ and HCO_3^- . Consequently, changes to any one of these molecules are reflected by variations in the other two, and all three (CO_2 , HCO_3^- , and pH) play central roles in biology. Cellular enzymes and chemical reactions are sensitive to pH, and cells actively transport H^+ and HCO_3^- across their cell membrane to maintain intracellular pH (pH_i) (reviewed in [1, 2]). Plus, $\text{CO}_2/\text{HCO}_3^-$ buffers intracellular and extracel-

lular physiological fluids and is the essential substrates or end products of biological calcification, photosynthesis, and respiration. Thus, cellular homeostasis and organismal homeostasis depend on sensing and tightly regulating levels of CO_2 , HCO_3^- , and H^+ .

While all cells must possess mechanisms to sense, and respond to, the levels of CO_2 , HCO_3^- , and H^+ , multicellular organisms have specialized cells which measure intracellular HCO_3^- ($[\text{HCO}_3^-]_i$) and pH_i as surrogates for the levels of CO_2 , pH (pH_e), and $[\text{HCO}_3^-]$ ($[\text{HCO}_3^-]_e$) in their immediate environment. In these sensory cells, variations in pH_i or $[\text{HCO}_3^-]_i$ elicit such diverse responses as secretion/absorption of H^+ , HCO_3^- , and other ions, as adjustments in lung ventilation rate, as changes in metabolism, and as regulation of gene expression.

Implicit in the definition of a biological sensor, in addition to “sensing” at least one of the relevant variables, a CO_2 , HCO_3^- , or pH sensor must trigger appropriate downstream responses. This signaling function can be achieved by altering membrane potential, by producing a second messenger molecule, or by modulating other proteins and enzymes via allosteric conformational changes or posttranslational modifications such as phosphorylation/dephosphorylation. In this review, we describe recent advances about how molecular pH, HCO_3^- , and/or CO_2 sensors function.

pH sensors

Extracellular pH (pH_e) sensors

Organisms sense pH_e to maintain and restore systemic acid/base balance, to adjust cellular metabolism to environmental conditions, and to detect and transduce sensory stimuli.

M. Tresguerres · J. Buck · L. R. Levin (✉)
Department of Pharmacology, Weill Cornell Medical College,
Cornell University,
New York, USA
e-mail: llevin@med.cornell.edu

J. Buck
e-mail: jobuck@med.cornell.edu

At least three G protein-coupled receptors (GPCRs) are activated by physiological increases in extracellular $[H^+]_e$ (decreases in pH_e): OGR1, GPR4, and TDAG8. In general, these pH_e -sensing GPCRs are inactive at $pH_e > 7.5$ and are fully activated at $pH_e < 6.8$. The mechanism of activation by H^+ probably involves the protonation of histidines exposed to the extracellular medium [3, 4], inducing conformational changes that promote activation of the particular downstream heterotrimeric G proteins. The end results are increased production of the intracellular second messengers cAMP (via G_s stimulation of transmembrane adenylyl cyclase activity) or IP_3 and diacylglycerol (via G_q family stimulation of phospholipase C activity).

Ovarian cancer G protein-coupled receptor 1 (OGR1, GPR68) stimulation of IP_3 production has been reported in osteoblasts [3] and osteoclasts [5], where it is thought to play roles in differentiation, metabolism, and systemic pH homeostasis [3, 5–8], as well as in aortic smooth muscle cells [6], where it is thought to mediate vasorelaxation via production of prostaglandin I_2 [7]. OGR1-dependent IP_3 formation is inactive at pH_e 7.8 and maximally active at pH_e 6.8 [3]. Activation of OGR1 also leads to the formation of cAMP in aortic smooth muscle cells [6]; however, this cAMP increase may be secondary to the IP_3 -elicited increase in intracellular calcium, rather than through direct G_s stimulation of a transmembrane adenylyl cyclase. Additionally, OGR1 is detectable in virtually every tissue tested and has been specifically identified in lung, kidney, and nervous system [3, 6, 8], suggesting there are additional roles of pH_e sensing.

GPR4 (aka GPRC6.1 [9]) is predominantly found in lung, although it is also present in kidney, heart, liver, skeletal muscle, ovary, and placenta [8–10]. In an over-expression system, GPR4 was linked to the activation of G_s and production of cAMP with pH responsiveness similar to OGR1 [3]; however, its signaling in more physiologically relevant contexts (i.e., aortic smooth muscle cells) remains unclear [7]. GPR4 is important for regulating angiogenesis [10], and it may also be important for the inflammatory response in endothelial cells [8].

The third H^+ -sensing GPCR is T cell death-associated gene (8TDAG8, GPR65), which seems to be exclusively present in immune cells and lymphoid tissues [6, 11]. Activation of TDAG8 by acidic pH_e leads the production of cAMP in a variety of transfected cultured cell lines [4, 12, 13] as well as in thymocytes and splenocytes [13]. TDAG8 overexpression in thymocytes promoted glucocorticoid-induced apoptosis [14]. TDAG8 genetic knockdown did not affect immune development or glucocorticoid-induced apoptosis [15]; however, the lack of phenotype may be due to compensation by other pH-sensing GPCRs or redundant pH-sensing mechanisms.

In addition to the H^+ -sensing GPCRs, pH_e is also sensed via H^+ -sensitive ion channels, for example, some inwardly

rectifying K^+ channels are regulated by intra- and/or extracellular pH. In general, acidic pH inhibits channel current while alkaline pH enhances it (reviewed in [16]). The better-characterized pH-sensitive K^+ channels are the rat outer medullary K^+ channel (ROMK) [17] and the TWIK-related acid-sensitive K^+ channel (TASK) [18]. ROMK, TASK, and related inwardly rectifying K^+ channels have been proposed to function as pH sensors throughout the body (especially in the kidney and in the brain stem) by modulating cell membrane potential (recently reviewed in [16, 19, 20]). Other H^+ -sensitive ion channels are predominantly found in sensory neurons and are involved in the signal transduction of pain or taste. These channels are gated by increases in $[H^+]_e$, and their activation stimulates conductance of K^+ , Na^+ , and/or Ca^{2+} , which triggers action potentials (Fig. 1). There are two major groups of H^+ -stimulated sensory channels: transient receptor potential (TRP) channels and acid-sensing ion channels (ASICs).

Polycystic kidney disease 2-like 1 (PKD2L1) is a TRP channel stimulated by low pH_e . PKD2L1 is present in taste receptor cells in the tongue, where it transduces sour stimuli (i.e., $pH_e < 5$) [21]. PKD2L1 is also present in a discrete population of neurons around the central canal of the spinal cord, which respond to minute pH variations around pH 7.4 [21]. This differential sensitivity to pH_e , as well as its functional expression on cellular membranes, is presumed to be due to tissue-specific association with modifier proteins [21, 22]. Additional information can be found in recent reviews on taste receptors [23] and PKD channels [24]. PKD2L1 is also involved in CO_2 sensing; in tongue taste buds, PKD2L1 senses extracellular acidification




Fig. 1 Extracellular pH/ CO_2 sensing by H^+ -gated channels in sensory neurons. *a* Extracellular H^+ , which could derive from elevated CO_2 (*b*), stimulate cation currents. Examples include TRP channels, such as PKD2L1 and TRPV1, and ASICs. If coexpressed with extracellular carbonic anhydrase IV (CA_{IV}), these channels can indirectly sense extracellular CO_2

subsequent to carbonic anhydrase IV-dependent hydration of CO_2 (see below).

A second TRP channel activated by acidic pH_e is the vanilloid receptor-1 (VR1 or TRPV1). At $\text{pH}_e < 5.9$, TRPV1 channels display increased open probability [25]. The stimulatory mechanism depends on the interaction of protons with two glutamic acid residues in the extracellular loop (reviewed in [26]). Because TRPV1 is also stimulated by capsaicin (the main ingredient of hot chili peppers) and heat [27], this channel is thought to sense burning pain throughout the body [28, 29].

Acid-sensing ion channels (ASICs), which are predominantly expressed in neuronal and neuroepithelial tissues [30–33], are activated by acidic $\text{pH}_e < 7$ [32]. ASICs, like all members of the degenerin family of channels, are more permeable to Na^+ , but they also transport Ca^{2+} and K^+ [31, 32]. There are several ASIC isoforms and splice variants, and their association into hetero- or homo-multimers may confer specific regulatory and kinetic properties in different tissues [33, 34].

In ASICs, gating by protons involves multiple residues in the extracellular domain, most notably carboxyl–carboxylate interactions between amino acids in the acidic pocket. These interactions result in the movement of a distal domain that blocks or unblocks the channel pore (ball and socket joint) [35]. Proton gating may also involve displacing Ca^{2+} ions which stabilize the closed state (reviewed in [36]).

ASICs are known to be important for nociception and taste [30–33], and they are also implicated in touch sensation in hairy skin [37], control of circulation by arterial baroreceptors [38], and pH_e sensing by carotid body glomus cells [39]. In addition, certain ASIC variants are present in the pituitary gland [40], bone [41], and smooth muscle [42] (among others [43]), with proposed roles in endocrine function, mineralization, and cell migration and control of circulation, respectively. However, it is not known if these functions are related to pH_e sensing. Finally, similar to PKD2L1, ASICs can also mediate CO_2 sensing via the associated variation in pH_e [44] (see below).

Intracellular pH (pH_i) sensors

Originally, it was posited that the H^+ and HCO_3^- transporting proteins which restore pH_i would be directly responsible for pH_i sensing. While it is true that a subset of these transporters are allosterically modulated by pH_i (e.g., sodium/hydrogen exchanger (NHE) [45]), their responsiveness is not sufficiently sensitive: pH_i regulation demands finer control than the observed kinetic responsiveness of NHEs and other pH_i regulatory ion transporting proteins (reviewed in [46]).

One “signaling” pH_i sensor is Pyk2, a member of the focal adhesion kinase family of tyrosine kinases [47]. Pyk2

is directly activated by acidic pH, both in cultured cells and in cell-free systems [48]. Although the exact molecular mechanism for acid activation is not known, it results in increased affinity for Pyk2’s substrate ATP (decreased K_m) without changing its V_{max} and is dependent on Ca^{2+} ions in the medium [48]. The initial steps of Pyk2 activation by acidic pH are autophosphorylation of tyrosine 402 [49] followed by a physical association with [48], and phosphorylation of, the tyrosine kinase c-Src [49]. While both c-Src and Pyk2 are activated by acidic pH [50], Pyk2 is considered the pH_i sensor because its activation precedes the other events, and c-Src activation is prevented by dominant-negative Pyk2 or siRNA-mediated inhibition of Pyk2 [48].


One of the consequences of acid stimulation of Pyk2 is NHE3 activation [48], which may be mediated by enhanced NHE3 transcription [51, 52] and/or by increasing NHE3 insertion into apical membranes [52]. NHE3 activation enhances H^+ secretion to restore pH_i (reviewed in [49]). The Pyk2/c-Src/NHE3 pathway is thought to be important for the compensation of systemic metabolic acidosis by the renal proximal tubule (reviewed in [46, 49]) (Fig. 2). Pyk2 and c-Src also mediate the activation of the $\text{Na}^+/\text{HCO}_3^-$ cotransporter 1 (NBC1, Slc4a4) in response to intracellular acidification by CO_2 [53]. Thus, Pyk2-mediated responses to acidosis can involve increases in both apical H^+ secretion via NHE3 and basolateral HCO_3^- absorption via NBC1 (Fig. 2). Finally, Pyk2 and c-Src are also important for the normal functioning of osteoclasts [54] and endothelial cells [55, 56], suggesting their pH_i sensory role extends beyond the kidney (reviewed in [46]).

The renal proximal tubule possesses a pH_i -sensing pathway independent of Pyk2 which leads to ERK activation via an unknown mechanism [49, 57]. One candidate for this Pyk2-independent pH_i sensing would be HCO_3^- -sensitive soluble adenylyl cyclase (sAC) (described below). The sAC has been shown to be upstream of ERK activation in other cells [58], and it is present in the renal proximal tubule [59]. However, the involvement of sAC in renal ERK activation has not yet been examined.

The pH-sensing protein kinases are also implicated in bacteria. *Salmonella* sense the lower pH inside the mammalian phagosome to stimulate transcription of the genes essential for virulence. In this system, the sensor for lowered pH is kinase PhoQ, which is directly activated by exposure of the sensor domain to pH 5.5 [60]. Also in *Salmonella*, pH_i sensing is important for activation of the machinery necessary for delivering toxin to their host’s cells, but in this case, the sensor is not yet known [61].

Bacteria also sense pH_i via second messenger generation. The adenylyl cyclase from *Mycobacterium tuberculosis* (Rv1264) [62] is activated by low pH. The molecular mechanism that confers Rv1264 pH sensitivity depends on

Fig. 2 Intracellular pH sensing by the tyrosine kinase Pyk2 in the renal proximal tubule. *a* Intracellular H^+ activate Pyk2 and c-Src, which promote apical H^+ secretion and Na^+ absorption by *b* sodium hydrogen exchanger 3 (NHE3) and *c* Na^+ and HCO_3^- absorption by sodium bicarbonate cotransporter (NBC)



the separation of an inhibitory subunit from the catalytic subunit [62]. Although Rv1264 was hypothesized to be important for sensing and counteracting acidification of phagolysosomes during host invasion [62], mutants lacking this cyclase demonstrated normal virulence [63]; no other *in vivo* role of Rv1264 is known.

The adenylyl cyclases from enteric bacteria (CyaA) such as *Escherichia coli* are also pH sensitive, but they are activated by high pH [64]. Purified CyaA demonstrates a very steep pH-dependency in the physiological range (between pH 7 and 9) *in vitro*, and while CyaA mediates the well-studied process of catabolite repression (reviewed in [65]), there is no defined role for pH_i sensing in this process. Conversely, it has been postulated that cAMP mediates physiological responses to pH_i changes [66, 67], but a definitive role for CyaA has not been established. Even though orthologs of Rv1264 and CyaA have not been found in eukaryotes, these studies reveal an evolutionarily ancient connection between second messenger signaling and pH_i sensing, and establish the paradigm of nucleotidyl cyclases acting as molecular sensors of pH_i .

Recent reviews about other molecular pH_i -sensing mechanisms can be found in [68] and [69].

Bicarbonate (HCO_3^-) sensors

Inside cells, HCO_3^- is used as a cofactor for carboxylation reactions, particularly via the vitamin K-dependent carboxylases, whose substrates play roles in blood coagulation, apoptosis, bone mineralization, calcium homeostasis, growth control, and signal transduction in the brain [70, 71]. However, vitamin K-dependent carboxylase is maximally stimulated at a $[HCO_3^-] < 2$ mM [72], which is way below its

intracellular level, indicating that their activity would not fluctuate with physiologically relevant changes in bicarbonate. Thus, although they are HCO_3^- -utilizing enzymes involved in signaling cascades, they do not satisfy the requirements for functioning as HCO_3^- sensors.

In contrast, two vertebrate signaling enzymes are responsive to physiologically relevant changes in intracellular $[HCO_3^-]$ —soluble adenylyl cyclase (sAC, also known as ADCY10 and Sacy) and guanylyl cyclase-D (GC-D) [73–75]—for example, sAC has been demonstrated to be modulated *in vivo* by $[HCO_3^-]$ which (a) originates via carbonic anhydrase-dependent hydration of exogenous CO_2 [59, 75–77], (b) enters the cell via DIDS-sensitive NBCs [Choi et al., unpublished], and (c) is generated from metabolic CO_2 production inside mitochondria [78, 79] (Fig. 3).

sAC is directly activated by HCO_3^- to produce the second messenger cAMP [73]. sAC is widely expressed [80–82], and it is distributed throughout the cell's cytoplasm and in organelles such as the nucleus, centriole, mitotic spindle, mid-body, mitochondria [83], and in cilia [84]. sAC orthologs are present from cyanobacteria to mammals, suggesting that HCO_3^- sensing via sAC-like cyclases is evolutionarily conserved [73, 85].

A mechanistic model for cAMP formation from ATP by sAC-like enzymes has been deduced from the HCO_3^- -regulated cyanobacterial sAC homolog CyaC [86], which is both structurally and kinetically similar to mammalian sAC [86, 87]. Like other class III adenylyl cyclases, sAC requires two divalent cations for activity [88, 89]. Though active with Mg^{2+} as the only available divalent ion, addition of Ca^{2+} increases the affinity (decreases K_m) for its substrate ATP to values consistent with the concentration of ATP found inside the cells [90]. These data suggest




Fig. 3 Activation of sAC inside the cells. Production of cAMP by sAC can be modulated by HCO_3^- which *a* originates via carbonic anhydrase (CA) hydration of exogenous CO_2 , *b* enters the cell via sodium bicarbonate cotransporter (NBC), or *c* is generated from metabolic CO_2 production inside mitochondria

that in vivo mammalian sAC utilizes both Mg^{2+} and Ca^{2+} , and that its activity will be sensitive to ATP fluctuations inside the cells. The structure- and kinetics-based model predicts that Ca^{2+} bound to the γ -phosphate of ATP enters the catalytic site and coordinates with specific residues in the sAC catalytic center, resulting in an “open sAC state”. The second divalent metal, in this case a Mg^{2+} ion, then binds to the α -phosphate of ATP, leading to a “closed state” by interaction with a different set of catalytic residues. The change from “open” to “closed” states induces the release of the β - and γ -phosphates and esterification of the remaining alpha phosphate with C3 of the ribose in adenosine (“cyclizing”). HCO_3^- stimulates substrate turnover (V_{\max}) by inducing an allosteric change which leads to active site closure, recruitment of the catalytic Mg^{2+} , and rearrangement of the phosphates in the bound ATP, thus facilitating cAMP formation and release [87].

There is a single functional sAC gene in the human genome; however, it utilizes multiple promoters, and sAC mRNA undergoes extensive alternative splicing [80, 81, 84, 91]. Full-length mammalian sAC (sAC_f) is comprised of two heterologous catalytic domains (C1 and C2) which constitute the amino terminal 50 kDa of protein. The additional ~ 140 -kDa C terminus of sAC_f includes several putative regulatory domains such as an autoinhibitory region [92] and canonical P-loop and leucine zipper sequences [85] of as yet unknown functions. The minimal functional sAC variant, termed sAC_t , is a truncated form almost exclusively comprised of C1 and C2 [85, 91]. This

C1C2 protein has robust intrinsic basal and HCO_3^- -stimulated activities, and it is sensitive to all known sAC inhibitors. Other sAC variants have not yet been characterized in detail, but molecular studies predict the existence of C2-only isoforms resulting from alternative splicing and promoter utilization [80, 81, 84]. Existence of these C2 isoforms provide a rationale to explain why male infertility is the only reported phenotype of the existing sAC-knockout mouse model (which deleted exons encoding the C1 domain) [93–95]; this sAC-C1 knockout mouse seems to retain the alternative promoter generating the putative C2-only proteins [80]. Confirmation of this hypothesis awaits generation of KO mice which specifically disrupt the C2 domain.

For mammalian sAC, HCO_3^- half-maximal effect (EC_{50}) ranges between 10 and 25 mM [73, 90], which is roughly equivalent to the $[\text{HCO}_3^-]$ levels found inside the cells and in plasma. In contrast, the HCO_3^- EC_{50} for sAC from the dogfish shark is ~ 5 mM [76], reflecting the lower $[\text{HCO}_3^-]$ of the internal fluids of aquatic animals compared to air breathers. It is thus possible that the kinetics of HCO_3^- activation will be specific for each sAC ortholog so they reflect the characteristics of that species physiology.

sAC serves as a sensor of extracellular pH/ HCO_3^- in the acid/base regulatory epithelia of the mammalian epididymis [59, 96] and kidney collecting duct [97, 98], and in shark gills [76]. Despite the evolutionary and physiological diversity of these epithelia, the sAC-dependent mechanisms for sensing and regulating pH and $[\text{HCO}_3^-]$ in extracellular fluids are astonishingly similar (Fig. 4). In each case, the extracellular stimulus (i.e., elevated luminal pH or $[\text{HCO}_3^-]$) is transmitted inside of the cell either by HCO_3^- movement through anion transporters [99] or by entrance of CO_2 via diffusion (or by facilitated diffusion through aquaporins [76, 99, 100]). When CO_2 diffusion into the cell is the initiating signal, extracellular carbonic anhydrase IV presumably functions to rapidly equilibrate increases in $[\text{HCO}_3^-]_e$ into readily available CO_2 [77, 99], which can diffuse into the cell. A second carbonic anhydrase (type II) inside the cell would be essential for hydrating CO_2 back into H^+ and HCO_3^- [59, 77, 99].

Regardless of how the bicarbonate enters the cells, in these pH/ HCO_3^- sensing epithelia, the elevated $[\text{HCO}_3^-]_i$ stimulates sAC to produce cAMP, which activates protein kinase A (PKA) [96, 98]. The end effect is translocation of vacuolar proton pump (V-ATPase)-containing vesicles into the cell membrane facing the increase in $\text{pH}_e/[\text{HCO}_3^-]_e$. In the clear cells of the epididymis and in renal A-type cells, the sAC-dependent V-ATPase translocation is into the apical membrane, producing the elongation of apical microvilli [59, 98, 101], while in shark gill base-secreting cells, the sAC-dependent V-ATPase translocation is into the basolateral membrane [76, 77, 102, 103]. The transport of




Fig. 4 Sensing and regulation of systemic pH and acid/base status by carbonic anhydrase (CA), sAC, and vacuolar proton pump (VHA). Alkalosis due to elevated extracellular HCO_3^- and/or pH results in elevated extracellular CO_2 , a reaction catalyzed by extracellular CA_{IV} . CO_2 diffuses inside the cell, where it is hydrated into H^+ and HCO_3^- . sAC is activated by intracellular HCO_3^- to produce cAMP, which promotes (via PKA) the insertion of VHA-containing vesicles into the cell membrane facing the alkalosis. Membrane-inserted VHAs secrete H^+ , which counteract the alkalosis. sAC may also modulate the activity of HCO_3^- transporters in the opposite membrane. In the clear cells of the epididymis and in A-type renal intercalated cells, Side A is the apical (mucosal) side and Side B is the basolateral (serosal) side. The polarity is reversed in base-secreting cells of the shark gill and in B-type renal intercalated cells

H^+ to the side with the original alkalotic stimulus not only counteracts the original alkalosis but it also supplies the H^+ which can combine with the elevated HCO_3^- to generate the CO_2 which fuels the cycle. In the whole animal experiments in the shark, this cellular mechanism was demonstrated to be relevant for systemic pH regulation. The components of this self-regulating loop, sAC, carbonic anhydrase, and V-ATPase are evolutionarily ancient enzymes, suggesting that sensing and regulation of pH/ $[\text{HCO}_3^-]$ via this signaling module is likely to be conserved. Variations of this mechanism due to differential CO_2 permeability, cell polarization, and association with other sensors and/or downstream effector proteins could result in additional homeostatic mechanisms for $\text{CO}_2/\text{HCO}_3^-/\text{pH}$.

Bicarbonate sensing via sAC is not limited to systemic acid/base regulation. sAC activity was originally detected in mammalian testis [104, 105], specifically in male germ cells. A biochemically related but particulate activity was

detected in spermatozoa [104], and its activity was postulated to be downstream of a NaHCO_3 signal [106–109]. Once the sAC gene was cloned [85], it was confirmed that sAC isoforms are highly expressed in testis and sperm [82]. Prior to ejaculation, sperm are stored in the lumen of the epididymis, where $[\text{HCO}_3^-]$ is much lower than in plasma (≤ 5 mM) [110]. Sperm are immotile and incompetent to fertilize an egg while in the epididymis, but upon ejaculation, sperm are mixed with prostatic and seminal fluids containing 25 mM HCO_3^- , which enters sperm via $\text{Na}^+/\text{HCO}_3^-$ cotransporters [111]. The HCO_3^- influx induces two stages of sperm activation essential for fertilization, motility, and capacitation [112–114], which were pharmacologically [95] and genetically [93–95, 115] confirmed to be due to sAC. Thus, sAC functions as a physiological bicarbonate sensor in mammalian sperm essential for fertilization.

In recent years, HCO_3^- sensing via sAC has also been associated with the regulation of NaCl absorption in mammalian kidney [116] and fish intestine [117]; cystic fibrosis transmembrane conductance regulator (CFTR) in human airways [118, 119] and bovine cornea [120]; osteoclast formation [121], early embryo development [122], and K^+ secretion in colon [123]; and fluid secretion in cholangiocytes [124] (Table 1).

Like sAC, GC-D is directly activated by HCO_3^- , but GC-D produces the second messenger cGMP instead of cAMP [74, 75]. The EC_{50} of GC-D for $[\text{HCO}_3^-]$ is ~ 20 mM, both in intracellular cGMP accumulation experiments in cells overexpressing GC-D and in cGMP production assays with purified recombinant GC-D catalytic domain [74, 75]. In rodents, GC-D is present primarily, if not exclusively, in olfactory neurons [125], and it appears to be a pseudogene in primates [126]. The proposed function for GC-D is as a putative sensor of atmospheric CO_2 in rodents [75, 127]; however, it is difficult to rectify how an enzyme whose EC_{50} for $[\text{HCO}_3^-]$ corresponds to $\sim 4\%$ CO_2 could sense changes in atmospheric CO_2 ($\sim 0.03\%$).

Carbon dioxide (CO_2) sensors

In insects and nematodes, CO_2 sensing induces either attraction or avoidance depending on the species and the particular neurons involved (recently reviewed in [128]). In CO_2 -sensing insects, like the malaria-carrying *Anopheles* mosquito and the fruit fly *Drosophila*, gaseous CO_2 is sensed by chemosensing neurons of the olfactory system [129]. *Drosophila* also sense CO_2 in solution, but this is mediated by different neurons of the gustatory system [130]. In fruit flies and the mosquito, two seven-transmembrane spanning members of the gustatory receptor family (Gr21A and Gr63A) are essential for sensing volatile CO_2 [131];

Table 1 Putative CO₂/pH/HCO₃⁻ sensing via sAC

Function	
pH and [HCO ₃ ⁻] sensing	
Acid/base and pH regulation	Regulation of V-ATPase recycling/translocation to the cell membrane in clear cells of the epididymis [59, 96], A-type intercalated cells of mammalian kidney [97, 98, 101], and base-secreting cells in shark gill [76, 77].
Ion and fluid transport	NaCl absorption in renal collecting duct [116] and toadfish intestine [117], HCO ₃ ⁻ transport in epithelial airways [118], Cl ⁻ secretion in cornea [120], K ⁺ secretion in colon [123], and fluid and Cl ⁻ secretion in pancreas [124].
Regulation of gene transcription	CREB phosphorylation [154]; CFTR gene transcription [119].
Sperm function	Sperm motility, capacitation, and possibly acrosome reaction [93–95, 115, 155].
Brain metabolic coupling	Glycogen breakdown and lactate release by astrocytes to be used by neurons [Choi HB, Gordon GRJ, Zhou N, Tai C, Ryu JK, McLarnon JG, Levin LR, Buck J, MacVicar BA (2010) Metabolic communication between astrocytes and neurons via bicarbonate responsive soluble adenylyl cyclase. unpublished].
Development and cell differentiation	Osteoclastogenesis [121]; early embryo development [122].
CO ₂ sensing	
Cilia movement	Ciliary beat frequency in airway epithelia [84].
Metabolic regulation	Mitochondrial oxidative phosphorylation in response to CO ₂ from TCA cycle [78, 79].

however, the exact molecular mechanisms, including whether they sense CO₂, HCO₃⁻, or H⁺, and their downstream signaling cascade(s), remain unknown (reviewed in [128]).

In the nematode *Caenorhabditis elegans*, CO₂ induces an avoidance behavior mediated via cGMP [132, 133]. The avoidance response is not induced by acidic pH or elevated [HCO₃⁻] in media, leading the authors to suggest that the stimulus is a change in the concentration of H⁺, HCO₃⁻ and/or CO₂ inside sensing cells [132]. The receptor guanylyl cyclase DAF-11 [133] as well as the cyclic nucleotide-gated channel subunits TAX-2 and TAX-4 [132, 133] are essential for *C. elegans* avoidance behavior and establish the link to cGMP. However, DAF-11 does not seem to be present in the same neurons as TAX-2 and TAX-4 [134], which possibly indicates a complex system with more than one chemoreceptor located in several sensing neurons.

In mammals, CO₂ is sensed in kidney, airways, tongue, and peripheral and central chemoreceptors. Although some prokaryote adenylyl cyclases may be directly activated by CO₂ [135], a similar regulation has not yet been unambiguously identified in mammals. In part, this may be due to the difficulties associated with differentiating between direct effects of CO₂ from effects due to pH and HCO₃⁻. The use of out-of-equilibrium CO₂/HCO₃⁻ solutions [136] is one of the few ways in which this issue can be addressed. Using this approach, sensitivity to pharmacological inhibitors implicates tyrosine kinases in promoting acid secretion/bicarbonate reabsorption in response to basolateral (“blood”) CO₂ in kidney proximal tubules [137, 138]. Interestingly, these experiments conclude that the inducing signal is CO₂/HCO₃⁻ and not pH_i, which

suggest involvement of an as yet undefined CO₂/HCO₃⁻ sensor instead of being mediated by Pyk2 and c-Src, which are proposed to function in the proximal tubules to activate apical H⁺ secretion by NHE3 in response to acidosis [46, 48, 49].

Because of the rapid equilibration of CO₂, HCO₃⁻, and H⁺ by carbonic anhydrases, in multiple physiological systems, CO₂ is sensed by a pH sensor or a HCO₃⁻ sensor coupled with a carbonic anhydrase, for example, as mentioned above, the mechanism for extracellular CO₂ sensing in sour taste receptor cells depends on extracellular carbonic anhydrase IV and PKD2L1 [139], the polycystic-kidney-disease-like acid-selective ion channel [21, 22]. Carbonic anhydrase IV hydrates CO₂ into HCO₃⁻ and H⁺ almost instantaneously; the elevated [H⁺] opens PKD2L1 channels, which allow the entry of cations into the cell and elicit the neuronal response [139] (Fig. 1). Extracellular CO₂ is also sensed via related variations in pH_e by acid-sensing ion channels (ASICs)¹ in the amygdala, and this pathway is important in fear-induced behaviors [44].

In vertebrates, peripheral chemoreceptors, specifically the aortic and carotid bodies, sense changes in arterial CO₂ and pH while central chemoreceptors sense CO₂ and pH of cerebral spinal fluid, and both types of chemoreceptors regulate breathing frequency and tidal volume. The chemosensing neurons may sense CO₂, [HCO₃⁻], pH_e, pH_i, transmembrane pH gradient, or oxidative stress, with the ultimate response possibly depending upon a combination

¹ In this case, it is not known whether the mechanism relies on the presence of a carbonic anhydrase.

of multiple sensors [140]. Although the potential direct effect of CO_2/H^+ on ion channels has been extensively studied ([141–144], see previous section on H^+ sensors), no molecular chemosensor has been unambiguously identified. Bicarbonate-sensitive sAC represents an additional intriguing candidate which has not yet been directly tested. sAC is present in carotid body [145], and cAMP levels in carotid body are elevated during hypercapnia [145, 146], reviewed in [140]. sAC was hypothesized to sense intracellular increases in $[\text{HCO}_3^-]$ derived from hypercapnia in type I (glomus) carotid body cells and to augment Ca^{2+} influx through L-type Ca^{2+} channels via PKA [146]. This stimulation is independent of changes in pH_e or pH_i [146] and, like the CO_2 sensing mechanisms explained above, probably depends on a carbonic anhydrase (c.f.[147]). We have confirmed the presence of sAC in rat glomus cells by immunofluorescence (unpublished observations), which also express abundant intracellular carbonic anhydrase [148, 149]. Additionally, L-type Ca^{2+} channels are activated by elevated CO_2 , independently of pH_i , in rat locus coeruleus neurons [143], suggesting sAC could also be important in central chemosensing; however, direct functional evidence for the role of sAC as a molecular sensor of hypercapnia in the carotid body or in other peripheral and central chemoreceptors, is lacking.

It has already been demonstrated that bicarbonate-regulated adenylyl cyclases, in conjunction with carbonic anhydrases, can function as CO_2 sensors. In the pathological fungi *Candida albicans* and *Cryptococcus neoformans*, the adenylyl cyclases (Cyr1) are essential for pathological differentiation. In both cases, the ACs are directly regulated by HCO_3^- [150, 151], and deletion of either the cyclase or the carbonic anhydrase abrogates pathogenicity. These pathogenic fungi know when they are inside an infected host, and hence when they should undergo their pathogenic differentiation, by sensing the host's internal $[\text{CO}_2]$ (5%), which is 150-fold higher than the ambient $[\text{CO}_2]_e$ (0.03%).

In mammals, CO_2 chemosensing via sAC has been demonstrated in the regulation of ciliary beat frequency (CBF) in lung epithelial cells. CO_2 exposure increases CBF in cultured human lung epithelial cells differentiated at an air-liquid interface. This response is mediated by $\text{CO}_2/\text{HCO}_3^-$ -dependent stimulation of sAC localized to the cilia's axoneme [84]. Carbonic anhydrases are present in the apical area of ciliated cells of bronchiolar epithelium [152], but a direct link of carbonic anhydrases to CBF has not yet been established. Analogy with other CO_2 -sensing mechanisms suggests that carbonic anhydrases mediate the stimulation of sAC in cilia. Because CO_2 levels are higher in exhaled breath compared to inhaled air, the regulation of CBF by sAC may ensure the proper clearance of mucus from airway epithelia, and impairments in this regulation may be associated with airway diseases such as asthma [84, 153].

Carbonic anhydrase and sAC also sense metabolically generated CO_2 inside mitochondria. Mitochondria are the predominant source of CO_2 in eukaryotic cells, and sAC is present inside mitochondria [83], where it coordinates the rate of ATP production via oxidative phosphorylation with nutritional availability [78, 79]. Mitochondrial sAC activity is stimulated by Krebs Cycle-generated CO_2 in a carbonic anhydrase-dependent manner. $\text{CO}_2/\text{HCO}_3^-$ stimulation of sAC activates intramitochondrial PKA which phosphorylates complex IV of the electron transport chain, increasing its rate and capacity to handle electrons. Thus, a mitochondrial CO_2 -carbonic anhydrase-sAC pathway couples nutritional status to ATP production to ensure minimal generation of reactive oxygen species, and sAC senses variations of intracellular (metabolic) changes in $\text{CO}_2/\text{HCO}_3^-$ in addition to $[\text{HCO}_3^-]$; secondary to extracellular (blood, environment) CO_2 , HCO_3^- and pH changes.

Conclusions

Biological systems sense CO_2 , HCO_3^- , and pH via multiple different types of sensors, and while the specifics may vary, the fundamentals of sensing are evolutionarily conserved. HCO_3^- and pH are directly sensed by protein kinases and nucleotidyl cyclases from bacteria (e.g., CyaA, PhoQ) through vertebrates (e.g., sAC, Pyk2), and CO_2 is indirectly sensed by HCO_3^- or pH sensors coupled to carbonic anhydrases. Thus, mechanisms for sensing pH, $[\text{HCO}_3^-]$, and CO_2 are present in all domains of life illustrating their importance for organismal function.

Acknowledgements We thank Dr. Carsten Wagner (University of Zurich, Switzerland) for the insightful comments on the manuscript.

References

1. Roos A, Boron WF (1981) Intracellular pH. *Physiol Rev* 61:296–434
2. Casey JR, Grinstein S, Orlowski J (2010) Sensors and regulators of intracellular pH. *Nat Rev Mol Cell Biol* 11:50–61
3. Ludwig MG, Vanek M, Guerini D, Gasser JA, Jones CE, Junker U, Hofstetter H, Wolf RM, Seuwen K (2003) Proton-sensing G-protein-coupled receptors. *Nature* 425:93–98
4. Wang J-Q, Kon J, Mogi C, Tobo M, Damirin A, Sato K, Komachi M, Malchinkhuu E, Murata N, Kimura T, Kuwabara A, Wakamatsu K, Koizumi H, Uede T, Tsujimoto G, Kurose H, Sato T, Harada A, Misawa N, Tomura H, Okajima F (2004) TDAG8 is a proton-sensing and psychosine-sensitive G-protein-coupled receptor. *J Biol Chem* 279:45626–45633
5. Komarova SV, Pereverzev A, Shum JW, Sims SM, Dixon SJ (2005) Convergent signaling by acidosis and receptor activator of NF- κ B ligand (RANKL) on the calcium/calcineurin/NFAT pathway in osteoclasts. *Proc Natl Acad Sci USA* 102:2643–2648

6. Tomura H, Mogi C, Sato K, Okajima F (2005) Proton-sensing and lysolipid-sensitive G-protein-coupled receptors: a novel type of multi-functional receptors. *Cell Signal* 17:1466–1476
7. Tomura H, Wang J-Q, Komachi M, Damirin A, Mogi C, Tobo M, Kon J, Misawa N, Sato K, Okajima F (2005) Prostaglandin I₂ production and cAMP accumulation in response to acidic extracellular pH through OGR1 in human aortic smooth muscle cells. *J Biol Chem* 280:34458–34464
8. Seuwen K, Ludwig M-G, Wolf RM (2006) Receptors for protons or lipid messengers or both? *J Recept Signal Transduct Res* 26:599–610
9. An S, Tsai C, Goetzl EJ (1995) Cloning, sequencing and tissue distribution of two related G protein-coupled receptor candidates expressed prominently in human lung tissue. *FEBS Lett* 375:121–124
10. Yang LV, Radu CG, Roy M, Lee S, McLaughlin J, Teitell MA, Iruela-Arispe ML, Witte ON (2007) Vascular abnormalities in mice deficient for the G protein-coupled receptor GPR4 that functions as a pH sensor. *Mol Cell Biol* 27:1334–1347
11. Choi J-W, Lee SY, Choi Y (1996) Identification of a putative G protein-coupled receptor induced during activation-induced apoptosis of T cells. *Cell Immunol* 168:78–84
12. Ishii S, Kihara Y, Shimizu T (2005) Identification of T cell death-associated gene 8 (TDAG8) as a novel acid sensing G-protein-coupled receptor. *J Biol Chem* 280:9083–9087
13. Radu CG, Nijagal A, McLaughlin J, Wang L, Witte ON (2005) Differential proton sensitivity of related G protein-coupled receptors T cell death-associated gene 8 and G2A expressed in immune cells. *Proc Natl Acad Sci USA* 102:1632–1637
14. Tosa N, Murakami M, Jia WY, Yokoyama M, Masunaga T, Iwabuchi C, Inobe M, Iwabuchi K, Miyazaki T, Onoe K, Iwata M, Uede T (2003) Critical function of T cell death-associated gene 8 in glucocorticoid-induced thymocyte apoptosis. *Int Immunol* 15:741–749
15. Radu CG, Cheng D, Nijagal A, Riedinger M, McLaughlin J, Yang LV, Johnson J, Witte ON (2006) Normal immune development and glucocorticoid-induced thymocyte apoptosis in mice deficient for the T-cell death-associated gene 8 receptor. *Mol Cell Biol* 26:668–677
16. Hibino H, Inanobe A, Furutani K, Murakami S, Findlay I, Kurachi Y (2010) Inwardly rectifying potassium channels: their structure, function, and physiological roles. *Physiol Rev* 90:291–366
17. Ho K, Nichols CG, Lederer WJ, Lytton J, Vassilev PM, Kanazirska MV, Hebert SC (1993) Cloning and expression of an inwardly rectifying ATP-regulated potassium channel. *Nature* 362:31–38
18. Duprat F, Lesage F, Fink M, Reyes R, Heurteaux C, Lazdunski M (1997) TASK, a human background K⁺ channel to sense external pH variations near physiological pH. *EMBO J* 16:5464–5471
19. Hebert SC, Desir G, Giebisch G, Wang W (2005) Molecular diversity and regulation of renal potassium channels. *Physiol Rev* 85:319–371
20. Heitzmann D, Warth R (2008) Physiology and pathophysiology of potassium channels in gastrointestinal epithelia. *Physiol Rev* 88:1119–1182
21. Huang AL, Chen XK, Hoon MA, Chandrashekar J, Guo W, Trankner D, Ryba NJP, Zuker CS (2006) The cells and logic for mammalian sour taste detection. *Nature* 442:934–938
22. Ishimaru Y, Inada H, Kubota M, Zhuang H, Tominaga M, Matsunami H (2006) Transient receptor potential family members PKD1L3 and PKD2L1 form a candidate sour taste receptor. *Proc Natl Acad Sci* 103:12569–12574
23. Chandrashekar J, Hoon MA, Ryba NJP, Zuker CS (2006) The receptors and cells for mammalian taste. *Nature* 444:288–294
24. Delmas P (2005) Polycystins: polymodal receptor/ion-channel cellular sensors. *Pflügers Arch Eur J Physiol* 451:264–276
25. Tominaga M, Caterina MJ, Malmberg AB, Rosen TA, Gilbert H, Skinner K, Raumann BE, Basbaum AI, Julius D (1998) The cloned capsaicin receptor integrates multiple pain-producing stimuli. *Neuron* 21:531–543
26. Tominaga M, Tominaga T (2005) Structure and function of TRPV1. *Pflügers Arch Eur J Physiol* 451:143–150
27. Caterina MJ, Schumacher MA, Tominaga M, Rosen TA, Levine JD, Julius D (1997) The capsaicin receptor: a heat-activated ion channel in the pain pathway. *Nature* 389:816–824
28. Caterina MJ, Julius D (2001) The vanilloid receptor: a molecular gateway to the pain pathway. *Ann Rev Neurosci* 24:487–517
29. Geppetti P, Trevisani M (2004) Activation and sensitisation of the vanilloid receptor: role in gastrointestinal inflammation and function. *Br J Pharmacol* 141:1313–1320
30. Chandrashekar J, Kuhn C, Oka Y, Yarmolinsky DA, Hummler E, Ryba NJP, Zuker CS (2010) The cells and peripheral representation of sodium taste in mice. *Nature* 464:297–301
31. Kellenberger S, Schild L (2002) Epithelial sodium channel/degnerin family of ion channels: a variety of functions for a shared structure. *Physiol Rev* 82:735–767
32. Waldmann R, Champigny G, Bassilana F, Heurteaux C, Lazdunski M (1997) A proton-gated cation channel involved in acid-sensing. *Nature* 386:173–177
33. Waldmann R, Lazdunski M (1998) H⁺-gated cation channels: neuronal acid sensors in the NaC/DEG family of ion channels. *Curr Opin Neurobiol* 8:418–424
34. Krishtal O (2003) The ASICs: signaling molecules? Modulators? *Trends Neurosci* 26:477–483
35. Jasti J, Furukawa H, Gonzales EB, Goux E (2007) Structure of acid-sensing ion channel 1 at 1.9 Å resolution and low pH. *Nature* 449:316–323
36. Gründer S, Chen X (2010) Structure, function, and pharmacology of acid-sensing ion channels (ASICs): focus on ASIC1a. *Int J Physiol Pathophysiol Pharmacol* 2:73–94
37. Price MP, Lewin GR, McIlwraith SL, Cheng C, Xie J, Heppenstall PA, Stucky CL, Mannsfeldt AG, Brennan TJ, Drummond HA, Qiao J, Benson CJ, Tarr DE, Hrstka RF, Yang B, Williamson RA, Welsh MJ (2000) The mammalian sodium channel BNC1 is required for normal touch sensation. *Nature* 407:1007–1011
38. Lu Y, Ma X, Sabharwal R, Snitsarev V, Morgan D, Rahmouni K, Drummond HA, Whiteis CA, Costa V, Price M, Benson C, Welsh MJ, Chapleau MW, Abboud FM (2009) The ion channel ASIC2 is required for baroreceptor and autonomic control of the circulation. *Neuron* 64:885–897
39. Tan Z-Y, Lu Y, Whiteis CA, Benson CJ, Chapleau MW, Abboud FM (2007) Acid-sensing ion channels contribute to transduction of extracellular acidosis in rat carotid body glomus cells. *Circ Res* 101:1009–1019
40. Gründer S, Geissler H-S, Bässler E-L, Ruppertsberg JP (2000) A new member of acid-sensing ion channels from pituitary gland. *NeuroReport* 11:1607–1611
41. Jahr H, van Driel M, van Osch GJVM, Weinans H, van Leeuwen JPTM (2005) Identification of acid-sensing ion channels in bone. *Biochem Biophys Res Commun* 337:349–354
42. Grifoni SC, Jernigan NL, Hamilton G, Drummond HA (2008) ASIC proteins regulate smooth muscle cell migration. *Microvasc Res* 75:202–210
43. Drummond HA, Jernigan NL, Grifoni SC (2008) Sensing tension: epithelial sodium channel/acid-sensing ion channel proteins in cardiovascular homeostasis. *Hypertension* 51:1265–1271
44. Ziemann AE, Allen JE, Dahdaleh NS, Drobot II, Coryell MW, Wunsch AM, Lynch CM, Faraci FM, Howard Iii MA, Welsh MJ, Wemmie JA (2009) The amygdala is a chemosensor that detects carbon dioxide and acidosis to elicit fear behavior. *Cell* 139:1012–1021

45. Aronson PS, Nee J, Suhm MA (1982) Modifier role of internal H⁺ in activating the Na⁺-H⁺ exchanger in renal microvillus membrane vesicles. *Nature* 299:161–163
46. Gluck SL (2004) Acid sensing in renal epithelial cells. *J Clin Invest* 114:1696–1699
47. Lev S, Moreno H, Martinez R, Canoll P, Peles E, Musacchio JM, Plowman GD, Rudy B, Schlessinger J (1995) Protein-tyrosine kinase Pyk2 involved in Ca²⁺-induced regulation of ion-channel and map kinase functions. *Nature* 376:737–745
48. Li SY, Sato S, Yang XJ, Preisig PA, Alpern RJ (2004) Pyk2 activation is integral to acid stimulation of sodium/hydrogen exchanger 3. *J Clin Invest* 114:1782–1789
49. Preisig PA (2007) The acid-activated signaling pathway: starting with Pyk2 and ending with increased NHE3 activity. *Kidney Int* 72:1324–1329
50. Yamaji Y, Tsuganezawa H, Moe OW, Alpern RJ (1997) Intracellular acidosis activates c-Src. *Am J Physiol Cell Physiol* 272:C886–C893
51. Ambuhl PM, Amemiya M, Danczkay M, Lotscher M, Kaissling B, Moe OW, Preisig PA, Alpern RJ (1996) Chronic metabolic acidosis increases NHE3 protein abundance in rat kidney. *Am J Physiol Renal Physiol* 271:F917–F925
52. Yang XJ, Amemiya M, Peng Y, Moe OW, Preisig PA, Alpern RJ (2000) Acid incubation causes exocytic insertion of NHE3 in OKP cells. *Am J Physiol Cell Physiol* 279:C410–C419
53. Espiritu DJD, Bernardo AA, Robey RB, Arruda JAL (2002) A central role for Pyk2-Src interaction in coupling diverse stimuli to increased epithelial NBC activity. *Am J Physiol Renal Physiol* 283:F663–F670
54. Soriano P, Montgomery C, Geske R, Bradley A (1991) Targeted disruption of the C-Src protooncogene leads to osteopetrosis in mice. *Cell* 64:693–702
55. Orr AW, Murhpy-Ulrich JE (2004) Regulation of endothelial cell function by FAK and Pyk2. *Front Biosci* 9:1254–1266
56. Kodama H, Fukuda K, Takahashi E, Tahara S, Tomita Y, Ieda M, Kimura K, Owada KM, Vuori K, Ogawa S (2003) Selective involvement of p130Cas/Crk/Pyk2/c-Src in endothelin-1-induced JNK activation. *Hypertension* 41:1372–1379
57. Tsuganezawa H, Sato S, Yamaji Y, Preisig PA, Moe OW, Alpern RJ (2002) Role of c-SRC and ERK in acid-induced activation of NHE3. *Kidney Int* 62:41–50
58. Ramos LS, Zippin JH, Kamenetsky M, Buck J, Levin LR (2008) Glucose and GLP-1 stimulate cAMP production via distinct adenylyl cyclases in INS-1E insulinoma cells. *J Gen Physiol* 132:329–338
59. Pastor-Soler N, Beaulieu V, Litvin TN, Da Silva N, Chen Y, Brown D, Buck J, Levin LR, Breton S (2003) Bicarbonate-regulated adenylyl cyclase (sAC) is a sensor that regulates pH-dependent V-ATPase recycling. *J Biol Chem* 278:49523–49529
60. Prost LR, Daley ME, Le Sage V, Bader MW, Le Moual H, Klevit RE, Miller SI (2007) Activation of the bacterial sensor kinase PhoQ by acidic pH. *Mol Cell* 26:165–174
61. Yu X-J, McGourty K, Liu M, Unsworth KE, Holden DW (2010) pH sensing by intracellular *Salmonella* induces effector translocation. *Science* 328:1040–1043
62. Tews I, Findeisen F, Sinning I, Schultz A, Schultz JE, Linder JU (2005) The structure of a pH-sensing mycobacterial adenylyl cyclase holoenzyme. *Science* 308:1020–1023
63. Dittrich D, Keller C, Ehlers S, Schultz JE, Sander P (2006) Characterization of a *Mycobacterium tuberculosis* mutant deficient in pH-sensing adenylyl cyclase Rv1264. *Int J Med Microbiol* 296:563–566
64. Tao M, Lipmann F (1969) Isolation of adenylyl cyclase from *Escherichia coli*. *Proc Natl Acad Sci USA* 63:86–92
65. Botsford JL, Harman JG (1992) Cyclic AMP in prokaryotes. *Microbiol Rev* 56:100–122
66. Castanie-Cornet M-P, Penfound TA, Smith D, Elliott JF, Foster JW (1999) Control of acid resistance in *Escherichia coli*. *J Bacteriol* 181:3525–3535
67. Ma Z, Richard H, Foster JW (2003) pH-dependent modulation of cyclic AMP levels and GadW-dependent repression of RpoS affect synthesis of the GadX regulator and *Escherichia coli* acid resistance. *J Bacteriol* 185:6852–6859
68. Srivastava J, Barber DL, Jacobson MP (2007) Intracellular pH sensors: design principles and functional significance. *Physiology* 22:30–39
69. Brown D, Paunescu TG, Breton S, Marshansky V (2009) Regulation of the V-ATPase in kidney epithelial cells: dual role in acid-base homeostasis and vesicle trafficking. *J Exp Biol* 212:1762–1772
70. Hassel B (2000) Carboxylation and anaplerosis in neurons and glia. *Mol Neurobiol* 22:21–40
71. Berkner KL (2008) Vitamin K-dependent carboxylation. *Vitam Horm* 78:131–156
72. Sadowski JA, Esmon CT, Suttie JW (1976) Vitamin K-dependent carboxylase. Requirements of the rat liver microsomal enzyme system. *J Biol Chem* 251:2770–2776
73. Chen Y, Cann MJ, Litvin TN, Iourgenko V, Sinclair ML, Levin LR, Buck J (2000) Soluble adenylyl cyclase as an evolutionarily conserved bicarbonate sensor. *Science* 289:625–628
74. Guo D, Zhang JJ, Huang XY (2009) Stimulation of guanylyl cyclase-D by bicarbonate. *Biochemistry* 48:4417–4422
75. Sun L, Wang H, Hu J, Han J, Matsunami H, Luo M (2009) Guanylyl cyclase-D in the olfactory CO₂ neurons is activated by bicarbonate. *Proc Natl Acad Sci USA* 106:2041–2046
76. Tresguerres M, Parks SK, Salazar E, Levin LR, Goss GG, Buck J (2010) Bicarbonate-sensing soluble adenylyl cyclase is an essential sensor for acid/base homeostasis. *Proc Natl Acad Sci USA* 107:442–447
77. Tresguerres M, Parks SK, Wood CM, Goss GG (2007) V-H⁺-ATPase translocation during blood alkalosis in dogfish gills: interaction with carbonic anhydrase and involvement in the postfeeding alkaline tide. *Am J Physiol Reg Int Comp Physiol* 292:R2012–R2019
78. Acin-Perez R, Salazar E, Brosel S, Yang H, Schon EA, Manfredi G (2009) Modulation of mitochondrial protein phosphorylation by soluble adenylyl cyclase ameliorates cytochrome oxidase defects. *EMBO Mol Med* 1:392–406
79. Acin-Perez R, Salazar E, Kamenetsky M, Buck J, Levin LR, Manfredi G (2009) Cyclic AMP produced inside mitochondria regulates oxidative phosphorylation. *Cell Metab* 9:265–276
80. Farrell J, Ramos L, Tresguerres M, Kamenetsky M, Levin LR, Buck J (2008) Somatic "soluble" adenylyl cyclase isoforms are unaffected in Sacy^{tm1Lex}/Sacy^{tm1Lex} "knockout" mice. *PLoS ONE* 3:e3251
81. Geng W, Wang Z, Zhang J, Reed BY, Pak CY, Moe OW (2005) Cloning and characterization of the human soluble adenylyl cyclase. *Am J Physiol Cell Physiol* 288:C1305–C1316
82. Sinclair ML, Wang XY, Mattia M, Conti M, Buck J, Wolgemuth DJ, Levin LR (2000) Specific expression of soluble adenylyl cyclase in male germ cells. *Mol Reprod Dev* 56:6–11
83. Zippin JH, Chen Y, Nahirney P, Kamenetsky M, Wuttke MS, Fischman DA, Levin LR, Buck J (2003) Compartmentalization of bicarbonate-sensitive adenylyl cyclase in distinct signaling microdomains. *FASEB J* 17:82–84
84. Schmid A, Sutto Z, Nlend MC, Horvath G, Schmid N, Buck J, Levin LR, Conner GE, Fregien N, Salathe M (2007) Soluble adenylyl cyclase is localized to cilia and contributes to ciliary beat frequency regulation via production of cAMP. *J Gen Physiol* 130:99–109
85. Buck J, Sinclair ML, Schapal L, Cann MJ, Levin LR (1999) Cytosolic adenylyl cyclase defines a unique signaling molecule in mammals. *Proc Natl Acad Sci USA* 96:79–84

86. Steegborn C, Litvin TN, Hess KC, Capper AB, Taussig R, Buck J, Levin LR, Wu H (2005) A novel mechanism for adenyllyl cyclase inhibition from the crystal structure of its complex with catechol estrogen. *J Biol Chem* 280:31754–31759
87. Steegborn C, Litvin TN, Levin LR, Buck J, Wu H (2005) Bicarbonate activation of adenyllyl cyclase via promotion of catalytic active site closure and metal recruitment. *Nat Struct Mol Biol* 12:32–37
88. Linder J (2006) Class III adenyllyl cyclases: molecular mechanisms of catalysis and regulation. *Cell Mol Life Sci* 63:1736–1751
89. Linder JU (2008) Structure-function relationships in *Escherichia coli* adenylate cyclase. *Biochem J* 415:449–454
90. Litvin TN, Kamenetsky M, Zarifyan A, Buck J, Levin LR (2003) Kinetic properties of "soluble" adenyllyl cyclase. Synergism between calcium and bicarbonate. *J Biol Chem* 278:15922–15926
91. Jaiswal BS, Conti M (2001) Identification and functional analysis of splice variants of the germ cell soluble adenyllyl cyclase. *J Biol Chem* 276:31698–31708
92. Chaloupka JA, Bullock SA, Iourgenko V, Levin LR, Buck J (2006) Autoinhibitory regulation of soluble adenyllyl cyclase. *Mol Reprod Dev* 73:361–368
93. Xie F, Garcia MA, Carlson AE, Schuh SM, Babcock DF, Jaiswal BS, Gossen JA, Esposito G, van Duin M, Conti M (2006) Soluble adenyllyl cyclase (sAC) is indispensable for sperm function and fertilization. *Dev Biol* 296:353–362
94. Esposito G, Jaiswal BS, Xie F, Krajnc-Franken MA, Robben TJ, Strik AM, Kuil C, Philipsen RL, van Duin M, Conti M, Gossen JA (2004) Mice deficient for soluble adenyllyl cyclase are infertile because of a severe sperm-motility defect. *Proc Natl Acad Sci USA* 101:2993–2998
95. Hess KC, Jones BH, Marquez B, Chen Y, Ord TS, Kamenetsky M, Miyamoto C, Zippin JH, Kopf GS, Suarez SS, Levin LR, Williams CJ, Buck J, Moss SB (2005) The "soluble" adenyllyl cyclase in sperm mediates multiple signaling events required for fertilization. *Dev Cell* 9:249–259
96. Pastor-Soler NM, Hallows KR, Smolak C, Gong F, Brown D, Breton S (2008) Alkaline pH- and cAMP-induced V-ATPase membrane accumulation is mediated by protein kinase A in epididymal clear cells. *Am J Physiol Cell Physiol* 294:C488–C494
97. Paunescu TG, Da Silva N, Russo LM, McKee M, Lu HA, Breton S, Brown D (2008) Association of soluble adenyllyl cyclase with the V-ATPase in renal epithelial cells. *Am J Physiol Renal Physiol* 294:F130–F138
98. Gong F, Alzamora R, Smolak C, Li H, Naveed S, Neumann D, Hallows KR, Pastor-Soler NM (2010) Vacuolar H⁺-ATPase apical accumulation in kidney intercalated cells is regulated by PKA and AMP-activated protein kinase. *Am J Physiol-Renal Physiol* 298:F1162–F1169
99. Pastor-Soler N, Pietrement C, Breton S (2005) Role of acid/base transporters in the male reproductive tract and potential consequences of their malfunction. *Physiology* 20:417–428
100. Boron WF (2006) Acid-base transport by the renal proximal tubule. *J Am Soc Nephrol* 17:2368–2382
101. Paunescu TG, Ljubojevic M, Russo LM, Winter C, McLaughlin MM, Wagner CA, Breton S, Brown D (2010) cAMP stimulates apical V-ATPase accumulation, microvillar elongation, and proton extrusion in kidney collecting duct A-intercalated cells. *Am J Physiol-Renal Physiol* 298:F643–F654
102. Tresguerres M, Parks SK, Katoh F, Goss GG (2006) Microtubule-dependent relocation of branchial V-H⁺-ATPase to the basolateral membrane in the Pacific spiny dogfish (*Squalus acanthias*): a role in base secretion. *J Exp Biol* 209:599–609
103. Tresguerres M, Katoh F, Fenton H, Jasinska E, Goss GG (2005) Regulation of branchial V-H⁺-ATPase Na⁺/K⁺-ATPase and NHE2 in response to acid and base infusions in the Pacific spiny dogfish (*Squalus acanthias*). *J Exp Biol* 208:345–354
104. Braun T (1975) The effect of divalent cations on bovine spermatozoal adenylate cyclase activity. *J Cyclic Nucleotide Res* 1:271–281
105. Neer EJ (1978) Multiple forms of adenylate cyclase. *Adv Cyclic Nucleotide Res* 9:69–83
106. Garbers DL, Tubb DJ, Hyne RV (1982) A requirement of bicarbonate for Ca²⁺-induced elevations of cyclic AMP in guinea pig spermatozoa. *J Biol Chem* 257:8980–8984
107. Garty NB, Salomon Y (1987) Stimulation of partially purified adenylate cyclase from bull sperm by bicarbonate. *FEBS Lett* 218:148–152
108. Okamura N, Tajima Y, Soejima A, Masuda H, Sugita Y (1985) Sodium bicarbonate in seminal plasma stimulates the motility of mammalian spermatozoa through direct activation of adenylate cyclase. *J Biol Chem* 260:9699–9705
109. Visconti PE, Muschietti JP, Flawia MM, Tezon JG (1990) Bicarbonate dependence of cAMP accumulation induced by phorbol esters in hamster spermatozoa. *Biochim Biophys Acta* 1054:231–236
110. Levine N, Marsh DJ (1971) Micropuncture studies of electrochemical aspects of fluid and electrolyte transport in individual seminiferous tubules, epididymis and vas deferens in rats. *J Physiol* 213:557–570
111. Demarco IA, Espinosa F, Edwards J, Sosnik J, de la Vega-Beltran JL, Hockensmith JW, Kopf GS, Darszon A, Visconti PE (2003) Involvement of a Na⁺/HCO₃⁻ cotransporter in mouse sperm capacitation. *J Biol Chem* 278:7001–7009
112. Boatman DF, Robbins RS (1991) Bicarbonate: carbon-dioxide regulation of sperm capacitation, hyperactivated motility, and acrosome reactions. *Biol Reprod* 44:806–813
113. Lee MA, Storey BT (1986) Bicarbonate is essential for fertilization of mouse eggs: mouse sperm require it to undergo the acrosome reaction. *Biol Reprod* 34:349–356
114. Visconti PE, Galantino-Homer H, Moore GD, Bailey JL, Ning X, Fornes M, Kopf GS (1998) The molecular basis of sperm capacitation. *J Androl* 19:242–248
115. Schuh SM, Carlson AE, McKnight GS, Conti M, Hille B, Babcock DF (2006) Signaling pathways for modulation of mouse sperm motility by adenosine and catecholamine agonists. *Biol Reprod* 74:492–500
116. Hallows KR, Wang HM, Edinger RS, Butterworth MB, Oyster NM, Li H, Buck J, Levin LR, Johnson JP, Pastor-Soler NM (2009) Regulation of epithelial Na⁺ transport by soluble adenyllyl cyclase in kidney collecting duct cells. *J Biol Chem* 284:5774–5783
117. Tresguerres M, Levin LR, Buck J, Grosell M (2010) Modulation of NaCl absorption by [HCO₃⁻] in the marine teleost intestine is mediated by soluble adenyllyl cyclase. *Am J Physiol Regul Integr Comp Physiol* 299:62–71
118. Wang Y, Lam CS, Wu F, Wang W, Duan Y, Huang P (2005) Regulation of CFTR channels by HCO₃⁻-sensitive soluble adenyllyl cyclase in human airway epithelial cells. *Am J Physiol Cell Physiol* 289:C1145–C1151
119. Baudouin-Legros M, Hamdaoui N, Borot F, Fritsch J, Ollero M, Planelles G, Edelman A (2008) Control of basal CFTR gene expression by bicarbonate-sensitive adenyllyl cyclase in human pulmonary cells. *Cell Physiol Biochem* 21:075–086
120. Sun XC, Zhai CB, Cui M, Chen Y, Levin LR, Buck J, Bonanno JA (2003) HCO₃⁻-dependent soluble adenyllyl cyclase activates cystic fibrosis transmembrane conductance regulator in corneal endothelium. *Am J Physiol Cell Physiol* 284:C1114–C1122
121. Geng W, Hill K, Zerwekh JE, Kohler T, Müller R, Moe OW (2009) Inhibition of osteoclast formation and function by bicarbonate: role of soluble adenyllyl cyclase. *J Cell Physiol* 220:332–340
122. Chen MH, Chen H, Zhou Z, Ruan YC, Wong HY, Lu YC, Guo JH, Chung YW, Huang PB, Huang HF, Zhou WL, Chan HC (2010) Involvement of CFTR in oviductal HCO₃⁻ secretion and

- its effect on soluble adenylyl cyclase-dependent early embryo development. *Hum Reprod* 25:1744–1754
123. Halm ST, Zhang J, Halm DR (2010) β -adrenergic activation of electrogenic K^+ and Cl^- secretion in guinea pig distal colonic epithelium proceeds via separate cAMP signaling pathways. *Am J Physiol Gastrointest Liver Physiol* 299:81–95
 124. Strazzabosco M, Fiorotto R, Melero S, Glaser S, Francis H, Spirli C, Alpini G (2009) Differentially expressed adenylyl cyclase isoforms mediate secretory functions in cholangiocyte subpopulation. *Hepatology* 50:244–252
 125. Fülle HJ, Vassar R, Foster DC, Yang RB, Axel R, Garbers DL (1995) A receptor guanylyl cyclase expressed specifically in olfactory sensory neurons. *Proc Natl Acad Sci USA* 92:3571–3575
 126. Young JM, Waters H, Dong C, Fülle H-J, Liman ER (2007) Degeneration of the olfactory guanylyl cyclase D gene during primate evolution. *PLoS ONE* 2:e884
 127. Hu J, Zhong C, Ding C, Chi Q, Walz A, Mombaerts P, Matsunami H, Luo M (2007) Detection of near-atmospheric concentrations of CO_2 by an olfactory subsystem in the mouse. *Science* 317:953–957
 128. Sharabi K, Lecuona E, Helenius IT, Beitel GJ, Sznajder JJ, Gruenbaum Y (2009) Sensing, physiological effects and molecular response to elevated CO_2 levels in eukaryotes. *J Cell Mol Med* 13:4304–4318
 129. de Bruyne M, Foster K, Carlson JR (2001) Odor coding in the *Drosophila* antenna. *Neuron* 30:537–552
 130. Fischler W, Kong P, Marella S, Scott K (2007) The detection of carbonation by the *Drosophila* gustatory system. *Nature* 448:1054–1057
 131. Jones WD, Cayirlioglu P, Kadow IG, Vosshall LB (2007) Two chemosensory receptors together mediate carbon dioxide detection in *Drosophila*. *Nature* 445:86–90
 132. Bretscher AJ, Busch KE, de Bono M (2008) A carbon dioxide avoidance behavior is integrated with responses to ambient oxygen and food in *Caenorhabditis elegans*. *Proc Natl Acad Sci USA* 105:8044–8049
 133. Hallem EA, Sternberg PW (2008) Acute carbon dioxide avoidance in *Caenorhabditis elegans*. *Proc Natl Acad Sci USA* 105:8038–8043
 134. Birnby DA, Link EM, Vowels JJ, Tian H, Colacurcio PL, Thomas JH (2000) A transmembrane guanylyl cyclase (DAF-11) and Hsp90 (DAF-21) regulate a common set of chemosensory behaviors in *Caenorhabditis elegans*. *Genetics* 155:85–104
 135. Hammer A, Hodgson DRW, Cann MJ (2006) Regulation of prokaryotic adenylyl cyclases by CO_2 . *Biochem J* 396:215–218
 136. Zhao J, Hogan EM, Bevensee MO, Boron WF (1995) Out-of-equilibrium CO_2/HCO_3^- solutions and their use in characterizing a new K/HCO_3^- cotransporter. *Nature* 374:636–639
 137. Zhou Y, Bouyer P, Boron WF (2006) Role of a tyrosine kinase in the CO_2 -induced stimulation of HCO_3^- reabsorption by rabbit S2 proximal tubules. *Am J Physiol Renal Physiol* 291:F358–F367
 138. Zhou Y, Zhao J, Bouyer P, Boron WF (2005) Evidence from renal proximal tubules that HCO_3^- and solute reabsorption are acutely regulated not by pH but by basolateral HCO_3^- and CO_2 . *Proc Natl Acad Sci USA* 102:3875–3880
 139. Chandrashekar J, Yarmolinsky D, von Buchholtz L, Oka Y, Sly W, Ryba NJP, Zuker CS (2009) The taste of carbonation. *Science* 326:443–445
 140. Putnam RW, Filosa JA, Ritucci NA (2004) Cellular mechanisms involved in CO_2 and acid signaling in chemosensitive neurons. *Am J Physiol Cell Physiol* 287:C1493–C1526
 141. Wellner-Kienitz M-C, Shams H, Scheid P (1998) Contribution of Ca^{2+} -activated K^+ channels to central chemosensitivity in cultivated neurons of fetal rat medulla. *J Neurophysiol* 79:2885–2894
 142. Pineda J, Aghajanian GK (1997) Carbon dioxide regulates the tonic activity of locus coeruleus neurons by modulating a proton- and polyamine-sensitive inward rectifier potassium current. *Neuroscience* 77:723–743
 143. Filosa JA, Dean JB, Putnam RW (2002) Role of intracellular and extracellular pH in the chemosensitive response of rat locus coeruleus neurones. *J Physiol (Lond)* 541:493–509
 144. Trapp S, Aller MI, Wisden W, Gourine AV (2008) A role for TASK-1 (KCNK3) channels in the chemosensory control of breathing. *J Neurosci* 28:8844–8850
 145. Nunes AR, Monteiro EC, Johnson SM, Gauda EB (2009) Bicarbonate-regulated soluble adenylyl cyclase (sAC) mRNA expression and activity in peripheral chemoreceptors. *Adv Exp Med Biol* 648:235–241
 146. Summers BA, Overholt JL, Prabhakar NR (2002) CO_2 and pH independently modulate L-type Ca^{2+} current in rabbit carotid body glomus cells. *J Neurophysiol* 88:604–612
 147. Gonzalez C, Almaraz L, Obeso A, Rigual R (1994) Carotid-body chemoreceptors: from natural stimuli to sensory discharges. *Physiol Rev* 74:829–898
 148. Nurse CA (1990) Carbonic-anhydrase and neuronal enzymes in cultured glomus cells of the carotid-body of the rat. *Cell Tissue Res* 261:65–71
 149. Rigual R, Iniguez C, Carreres J, Gonzalez C (1985) Carbonic-anhydrase in the carotid-body and the carotid-sinus nerve. *Histochemistry* 82:577–580
 150. Mogensen EG, Janbon G, Chaloupka J, Steegborn C, Fu MS, Moyrand F, Klengel T, Pearson DS, Geeves MA, Buck J, Levin LR, Muhlschlegel FA (2006) *Cryptococcus neoformans* senses CO_2 through the carbonic anhydrase Can2 and the adenylyl cyclase Cac1. *Eukaryot Cell* 5:103–111
 151. Klengel T, Liang WJ, Chaloupka J, Ruoff C, Schroppel K, Naglik JR, Eckert SE, Mogensen EG, Haynes K, Tuite MF, Levin LR, Buck J, Muhlschlegel FA (2005) Fungal adenylyl cyclase integrates CO_2 sensing with cAMP signaling and virulence. *Curr Biol* 15:2021–2026
 152. Spicer SS, Ge Z-H, Tashian RE, Hazen-Martin DJ, Schulte BA (1990) Comparative distribution of carbonic anhydrase isozymes III and II in rodent tissues. *Am J Anat* 187:55–64
 153. Sutto Z, Conner GE, Salathe M (2004) Regulation of human airway ciliary beat frequency by intracellular pH. *J Physiol* 560:519–532
 154. Zippin JH, Farrell J, Huron D, Kamenetsky M, Hess KC, Fischman DA, Levin LR, Buck J (2004) Bicarbonate-responsive "soluble" adenylyl cyclase defines a nuclear cAMP microdomain. *J Cell Biol* 164:527–534
 155. Beltran C, Vacquier VD, Moy G, Chen Y, Buck J, Levin LR, Darszon A (2007) Particulate and soluble adenylyl cyclases participate in the sperm acrosome reaction. *Biochem Biophys Res Commun* 358:1128–1135